

2003 Health and Preventive Practices Status Report

from Missouri County-level Study



Crawford County**

	County Prevalence (%)*	Central Region Prevalence (%)*	State Prevalence (%)*
G eneral			
Health Status (Fair or Poor)	24.7 H	18.6	16.9
No Health Coverage	10.8	13.0	12.3
B ehavioral Risk Factors			
Physical Inactivity	36.1 H	23.7	23.0
Current Smoker	26.6	27.9	26.5
Overweight (BMI 25-29.9)	35.3	36.6	35.3
Obese (BMI ≥ 30)	26.4	23.6	23.4
D isease and Conditions			
High Blood Pressure	30.3	27.7	28.5
High Cholesterol (35 years or older)	37.3	35.8	37.3
Asthma	9.3	10.9	10.7
Diabetes	7.1	7.4	7.2
Activity Limitation	22.1	19.4	17.8
C ancer Screening			
<i>Women</i>			
Never had a Mammogram	14.3	12.9	13.4
No Mammogram or Clinical Breast Exam in Last Year:			
40-49 years of age	40.0	46.3	49.4
50-64 years of age	33.5	31.2	35.9
65 and older	58.2	54.3	47.4
Never had a Pap Test:			
18-69 years of age	2.4	3.0	5.1
70 and older	11.9	8.8	6.4
No Pap Test in Last Year (18-29)	11.6	20.2	25.8
No Pap Test in Last 3 Years (30-69)	18.5	12.5	13.8
<i>Men and Women 50 Years or Older</i>			
Blood Stool Test:			
Never had a Blood Stool Test	66.8	59.9	57.0
No Blood Stool Test in Last Year	88.3 H	82.4	77.5
Sigmoidoscopy/Colonoscopy			
Never had a Sigmoidoscopy or Colonoscopy	45.2	50.3	49.2
No Sigmoidoscopy or Colonoscopy in Past Five Years	55.2	60.4	57.0

*Prevalence rates are age adjusted to the 2000 U.S. standard population

■ indicates that the prevalence rate is not significantly above or below state prevalence

H indicates that the prevalence rate is significantly above the state prevalence

L indicates that the prevalence rate is significantly below the state prevalence

Unless otherwise noted respondents are 18 years of age or older

** The best indicators for Crawford county were obtained by combining data from Crawford and Washington counties.